

RESEARCH ARTICLE SUMMARY

POPULATION GENETICS


Genomic evidence for the Pleistocene and recent population history of Native Americans

Maanasa Raghavan *et al.**

INTRODUCTION: The consensus view on the peopling of the Americas is that ancestors of modern Native Americans entered the Americas from Siberia via the Bering Land Bridge and that this occurred at least ~14.6 thousand years ago (ka). However, the number and timing of migrations into the Americas remain controversial, with conflicting interpretations based on anatomical and genetic evidence.

RATIONALE: In this study, we address four major unresolved issues regarding the Pleistocene and recent population history of Native Americans: (i) the timing of their divergence from their ancestral group, (ii) the number of migrations into the Americas, (iii) whether there was ~15,000 years of isolation of ancestral Native Americans in Beringia (Beringian

Incubation Model), and (iv) whether there was post-Pleistocene survival of relict populations in the Americas related to Australo-Melanesians, as suggested by apparent differences in cranial morphologies between some early ("Paleoamerican") remains and those of more recent Native Americans. We generated 31 high-coverage modern genomes from the Americas, Siberia, and Oceania; 23 ancient genomic sequences from the Americas dating between ~0.2 and 6 ka; and SNP chip genotype data from 79 present-day individuals belonging to 28 populations from the Americas and Siberia. The above data sets were analyzed together with published modern and ancient genomic data from worldwide populations, after masking some present-day Native Americans for recent European admixture.



Population history of present-day Native Americans. The ancestors of all Native Americans entered the Americas as a single migration wave from Siberia (purple) no earlier than ~23 ka, separate from the Inuit (green), and diversified into "northern" and "southern" Native American branches ~13 ka. There is evidence of post-divergence gene flow between some Native Americans and groups related to East Asians/Inuit and Australo-Melanesians (yellow).

RESULTS: Using three different methods, we determined the divergence time for all Native Americans (Athabascans and Amerindians) from their Siberian ancestors to be ~20 ka, and no earlier than ~23 ka. Furthermore, we dated the divergence between Athabascans (northern Native American branch, together

ON OUR WEB SITE

Read the full article at <http://dx.doi.org/10.1126/science.aab3884>

with northern North American Amerindians) and southern North Americans and South and Central Americans (southern Native American branch) to be ~13 ka. Similar divergence times from East Asian populations and a divergence time between the two branches that is close in age to the earliest well-established archaeological sites in the Americas suggest that the split between the branches occurred within the Americas. We additionally found that several sequenced Holocene individuals from the Americas are related to present-day populations from the same geographical regions, implying genetic continuity of ancient and modern populations in some parts of the Americas over at least the past 8500 years. Moreover, our results suggest that there has been gene flow between some Native Americans from both North and South America and groups related to East Asians and Australo-Melanesians, the latter possibly through an East Asian route that might have included ancestors of modern Aleutian Islanders. Last, using both genomic and morphometric analyses, we found that historical Native American groups such as the Pericúes and Fuego-Patagonians were not "relicts" of Paleoamericans, and hence, our results do not support an early migration of populations directly related to Australo-Melanesians into the Americas.

CONCLUSION: Our results provide an upper bound of ~23 ka on the initial divergence of ancestral Native Americans from their East Asian ancestors, followed by a short isolation period of no more than ~8000 years, and subsequent entrance and spread across the Americas. The data presented are consistent with a single-migration model for all Native Americans, with later gene flow from sources related to East Asians and, indirectly, Australo-Melanesians. The single wave diversified ~13 ka, likely within the Americas, giving rise to the northern and southern branches of present-day Native Americans. ■

The list of author affiliations is available in the full article online.

*Corresponding authors. E-mail: ewillierslev@snm.ku.dk (E.W.); rasmus_nielsen@berkeley.edu (R.N.); yss@berkeley.edu (Y.S.S.)

Cite this article as M. Raghavan *et al.*, *Science* 349, aab3884 (2015). DOI: 10.1126/science.aab3884

RESEARCH ARTICLE

POPULATION GENETICS

Genomic evidence for the Pleistocene and recent population history of Native Americans

Maanasa Raghavan,^{1*} Matthias Steinrücken,^{2,3,4*} Kelley Harris,^{5*} Stephan Schiffels,^{6*} Simon Rasmussen,^{7*} Michael DeGiorgio,^{8*} Anders Albrechtsen,^{9*} Cristina Valdiosera,^{1,10*} María C. Ávila-Arcos,^{1,11*} Anna-Sapfo Malaspinas,^{1*} Anders Eriksson,^{12,13} Ida Moltke,⁹ Mait Metspalu,^{14,15} Julian R. Homburger,¹¹ Jeff Wall,¹⁶ Omar E. Cornejo,¹⁷ J. Víctor Moreno-Mayar,¹ Thorfinn S. Korneliussen,¹ Tracey Pierre,¹ Morten Rasmussen,^{1,11} Paula F. Campos,^{1,18} Peter de Barros Damgaard,¹ Morten E. Allentoft,¹ John Lindo,¹⁹ Ene Metspalu,^{14,15} Ricardo Rodríguez-Varela,²⁰ Josefina Mansilla,²¹ Celeste Henrickson,²² Andaine Seguin-Orlando,¹ Helena Malmström,²³ Thomas Stafford Jr.,^{1,24} Suyash S. Shringarpure,¹¹ Andrés Moreno-Estrada,^{11,25} Monika Karmin,^{14,15} Kristiina Tambets,¹⁴ Anders Bergström,⁶ Yali Xue,⁶ Vera Warmuth,^{26,27} Andrew D. Friend,²⁸ Joy Singarayer,²⁹ Paul Valdes,³⁰ Francois Balloux,²⁶ Ilán Lebreiro,²¹ Jose Luis Vera,³¹ Hector Rangel-Villalobos,³² Davide Pettener,³³ Donata Luiselli,³³ Loren G. Davis,³⁴ Evelyn Heyer,³⁵ Christoph P. E. Zollikofer,³⁶ Marcia S. Ponce de León,³⁶ Colin I. Smith,¹⁰ Vaughan Grimes,^{37,38} Kelly-Anne Pike,³⁷ Michael Deal,³⁷ Benjamin T. Fuller,³⁹ Bernardo Arriaza,⁴⁰ Vivien Standen,⁴¹ Maria F. Luz,⁴² Francois Ricaut,⁴³ Niede Guidon,⁴² Ludmila Osipova,^{44,45} Mikhail I. Voevoda,^{44,46,47} Olga L. Posukh,^{44,45} Oleg Balanovsky,^{48,49} Maria Lavryashina,⁵⁰ Yuri Bogunov,⁴⁸ Elza Khusnutdinova,^{51,52} Marina Gubina,⁴⁴ Elena Balanovska,⁴⁹ Sardana Fedorova,^{53,54} Sergey Litvinov,^{14,51} Boris Malyarchuk,⁵⁵ Miroslava Derenko,⁵⁵ M. J. Mosher,⁵⁶ David Archer,⁵⁷ Jerome Cybulski,^{58,59,60} Barbara Petzelt,⁶¹ Joycelynn Mitchell,⁶¹ Rosita Worl,⁶² Paul J. Norman,⁶³ Peter Parham,⁶³ Brian M. Kemp,^{17,64} Toomas Kivisild,^{14,65} Chris Tyler-Smith,⁶ Manjinder S. Sandhu,^{6, 66} Michael Crawford,⁶⁷ Richard Villems,^{14,15} David Glenn Smith,⁶⁸ Michael R. Waters,^{69,70,71} Ted Goebel,⁶⁹ John R. Johnson,⁷² Ripan S. Malhi,^{19,73} Mattias Jakobsson,²³ David J. Meltzer,^{1,74} Andrea Manica,¹² Richard Durbin,⁶ Carlos D. Bustamante,¹¹ Yun S. Song,^{2,3,75†} Rasmus Nielsen,^{75†} Eske Willerslev^{1†}

How and when the Americas were populated remains contentious. Using ancient and modern genome-wide data, we found that the ancestors of all present-day Native Americans, including Athabascans and Amerindians, entered the Americas as a single migration wave from Siberia no earlier than 23 thousand years ago (ka) and after no more than an 8000-year isolation period in Beringia. After their arrival to the Americas, ancestral Native Americans diversified into two basal genetic branches around 13 ka, one that is now dispersed across North and South America and the other restricted to North America. Subsequent gene flow resulted in some Native Americans sharing ancestry with present-day East Asians (including Siberians) and, more distantly, Australo-Melanesians. Putative “Paleoamerican” relict populations, including the historical Mexican Pericúes and South American Fuego-Patagonians, are not directly related to modern Australo-Melanesians as suggested by the Paleoamerican Model.

It is generally agreed that ancestral Native Americans are descendants of Siberian peoples who traversed the Bering Land Bridge (Beringia) from northeast Asia in Late Pleistocene times, and although consensus has yet to be reached, it is mostly conceded that the Clovis archaeological complex, dating to ~13 thousands years ago (ka), does not represent the first migration as long supposed (1–7). Archaeological evidence accumulated over the past two decades indicates that people were south of the North

American continental ice sheets more than a millennium earlier and had reached as far south as southern South America by at least ~14.6 ka (1–3). Interpretations differ, however, regarding the precise spatiotemporal dynamics of the peopling process, owing to archaeological claims for a substantially earlier human presence predating the Last Glacial Maximum (LGM) (~20 ka) (8–10) and conflicting interpretations of the number and timing of migrations from Beringia based on anatomical and genetic evidence (11–16). Much

of the genetic evidence is from studies of mitochondrial DNA (mtDNA) and Y chromosome, which as single, uniparentally inherited loci are particularly subject to genetic drift and sex-biased demographic and cultural practices.

Among the principal issues still to be resolved regarding the Pleistocene and recent population history of Native Americans are (i) the timing of their divergence from their Eurasian ancestors; (ii) whether the peopling was in a single wave or multiple waves and, consequently, whether the genetic differences seen between major subgroups of Native Americans (such as Amerindian and Athabascan) result from different migrations or in situ diversification in the Americas (5, 6, 17, 18); (iii) whether the migration involved ~15,000 years of isolation in the Bering Strait region, as proposed by the Beringian Incubation Model to explain the high frequency of distinct and widespread American mitogenomes and private genetic variants (19–22); and last, (iv) whether there was post-divergence gene flow from Eurasia and possibly even population replacement in the Americas, the latter suggested by the apparent differences in skull morphology between some early (“Paleoamerican”) remains and those of more recent Native Americans (23–27). We address these issues using genomic data derived from modern populations, supplemented by ancient specimens that provide chronologically controlled snapshots of the genetics of the peopling process as it unfolded.

We sequenced 31 genomes from present-day individuals from the Americas, Siberia, and Oceania to an average depth of ~20×: Siberians—Altai ($n = 2$), Buryat ($n = 2$), Ket ($n = 2$), Koryak ($n = 2$), Sakha ($n = 2$), and Siberian Yupik ($n = 2$); North American Native Americans—Tsimshian ($n = 1$); southern North American and Central and South American Natives—Pima ($n = 1$), Huichol ($n = 1$), Aymara ($n = 1$), and Yukpa ($n = 1$); and Oceanians—Papuan ($n = 14$) (table S1) (28). All the genome-sequenced present-day individuals were previously genotyped by using single-nucleotide polymorphism (SNP) chips (4, 29–35) except for the Aymara individual, which was SNP chip-genotyped in this study (tables S3 and S4). They were selected on the basis of their ancestry profiles obtained with ADMIXTURE (36) so as to best represent their respective populations and to minimize recent genetic admixture from populations of western Eurasian origin (28). For populations represented by more than one individual, we also verified from the genotype data that the sequenced individuals did not represent close relatives (28). We additionally sequenced 23 genomes from ancient individuals dating between ~0.2 and 6 ka from North and South America, with an average depth ranging between 0.003× and 1.7×, including specimens affiliated to putative relict Paleoamerican groups such as the Pericúes from Mexico and Fuego-Patagonians from the southernmost tip of South America (table S5) (23, 26–28). Last, we generated SNP chip genotype data from 79 present-day individuals belonging to 28 populations from the Americas and Siberia (table S4) (28). All the

forementioned data sets were analyzed together with previously published genomes and SNP chip genotype data (tables S1, S3, and S4), masking the data for recent European admixture in some present-day Native American populations (28).

The structure of Native American populations and the timing of their initial divergence

We explored the genetic structure of Native American populations in the context of worldwide populations using ADMIXTURE (36), using a reference panel consisting of 3053 individuals from 169 populations (table S3) (28). The panel included SNP chip genotype data from present-day individuals generated in this study and previously published studies, as well as the 4000-year-old Saqqaq individual from Greenland (29) and the 12,600-year-old Anzick-1 (Clovis culture) individual from Montana (table S3) (5). When assuming the number of ancestral populations (K) to be four ($K = 4$), we found a Native American-specific genetic component, indicating a shared genetic ancestry for all Native Americans, including Amerindians and Athabascans (fig. S4). Assuming $K = 15$, there is structure within the Native Americans. Athabascans and northern

Amerindians (primarily from Canada) differ from the rest of the Native Americans in sharing their own genetic component (fig. S4). As reported previously, Anzick-1 falls within the genetic variation of southern Native Americans (5), whereas the Saqqaq individual shares genetic components with Siberian populations (fig. S4) (29).

To ascertain the population history of present-day Native American populations in relation to worldwide populations, we generated admixture graphs with TreeMix (28, 37). All of the modern Siberian and Native American genomes sequenced in this study, except for the North American Tsimshian genome that showed evidence of recent western Eurasian admixture (28), were used for this analysis, together with previously published genomes from Africa (Yoruba) (38), Europe (Sardinian and French) (38), East Asia (Dai and Han) (38), Siberia (Nivkh) (39) and the Americas (Karitiana, Athabaskan, and Greenlandic Inuit) (table S1) (5, 38, 39). The ancient individuals included in the analysis were Saqqaq, Anzick-1, and the 24,000-year-old Mal'ta child from south-central Siberia (4). By use of TreeMix, we affirmed that all Native Americans form a monophyletic group across all 10 migration parameter values, with further diversification into two branches, one

representing Amerindians (represented in this analysis by Amerindians from southern North America and Central and South America) and the other representing Athabascans (Fig. 1B and fig. S5). Paleo-Eskimos and Inuit were supported as a separate clade relative to the Native Americans, as reported previously (Fig. 1B and fig. S5) (29, 39). Our results show that the Siberian Yupik and Koryak are the closest Eurasian populations to the Americas, with the Yupik likely representing back-migration of the Inuit into Siberia (Fig. 1B and fig. S5).

To assess the pattern of the earliest human dispersal into the Americas, we estimated the timing of the divergence of ancestral Native Americans from East Asians (hereafter, including Siberians) using multiple methods. There is still some debate regarding mutation rates in the human genome (40), and this uncertainty could affect our estimates and results.

We applied diCal2.0 (method 1) (28), a new version of diCal (41) extended to handle complex demographic models involving multiple populations with migration (42), and an identity-by-state (IBS) tract method (method 2) (43) (supplementary materials, materials and methods 2) to the modern genomes data set (28). With these, we

¹Centre for GeoGenetics, Natural History Museum of Denmark, University of Copenhagen, Øster Voldgade 5–7, 1350 Copenhagen, Denmark. ²Computer Science Division, University of California, Berkeley, Berkeley, CA 94720, USA. ³Department of Statistics, University of California, Berkeley, Berkeley, CA 94720, USA. ⁴Department of Biostatistics and Epidemiology, University of Massachusetts, Amherst, MA 01003, USA. ⁵Department of Mathematics, University of California, Berkeley, Berkeley, CA 94720, USA. ⁶Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton CB10 1SA, UK. ⁷Center for Biological Sequence Analysis, Department of Systems Biology, Technical University of Denmark, Kemitorvet, Building 208, 2800 Kongens Lyngby, Denmark. ⁸Departments of Biology and Statistics, Pennsylvania State University, 502 Wartik Laboratory, University Park, PA 16802, USA. ⁹The Bioinformatics Centre, Department of Biology, University of Copenhagen, Ole Maaløes Vej 5, 2200 Copenhagen, Denmark. ¹⁰Department of Archaeology and History, La Trobe University, Melbourne, Victoria 3086, Australia. ¹¹Department of Genetics, School of Medicine, Stanford University, 300 Pasteur Drive, Lane Building, Room L331, Stanford, CA 94305, USA. ¹²Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK. ¹³Integrative Systems Biology Laboratory, King Abdullah University of Science and Technology (KAUST), Thuwal, 23955-6900, Kingdom of Saudi Arabia. ¹⁴Estonian Biocentre, Evolutionary Biology Group, Tartu 51010, Estonia. ¹⁵Department of Evolutionary Biology, University of Tartu, Tartu 51010, Estonia. ¹⁶Institute for Human Genetics, University of California San Francisco, 513 Parnassus Avenue, San Francisco, CA 94143, USA. ¹⁷School of Biological Sciences, Washington State University, Post Office Box 644236, Heald 429, Pullman, WA 99164, USA. ¹⁸Centro de Investigación en Ciencias del Mar y Limnología/Centro Interdisciplinar de Investigación Marina e Ambiental, Centro Interdisciplinar de Investigación Marina e Ambiental, Universidade do Porto, Rua dos Bragas 289, 4050-123 Porto, Portugal. ¹⁹Department of Anthropology, University of Illinois at Urbana-Champaign, 607 S. Mathews Avenue, Urbana, IL 61801, USA. ²⁰Centro Mixto, Universidad Complutense de Madrid-Instituto de Salud Carlos III de Evolución y Comportamiento Humano, Madrid, Spain. ²¹Instituto Nacional de Antropología e Historia, Moneda 13, Centro, Cuauhtémoc, 06060 Mexico City, Mexico. ²²University of Utah, Department of Anthropology, 270 S 1400 E, Salt Lake City, UT 84112, USA. ²³Department of Evolutionary Biology and Science for Life Laboratory, Uppsala University, Norbyvägen 18D, SE-752 36 Uppsala, Sweden. ²⁴Acceleration Mass Spectrometry 14C Dating Centre, Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, 8000 Aarhus, Denmark. ²⁵Laboratorio Nacional de Genómica para la Biodiversidad (LANGEBIO), Centro de Investigación y de Estudios Avanzados, Irapuato, Guanajuato 36821, Mexico. ²⁶Genetics Institute, University College London, Gower Street, London WC1E 6BT, UK. ²⁷Evolutionärsbiologiskt Centrum, Norbyvägen 18D, 75236 Uppsala, Sweden. ²⁸Department of Geography, University of Cambridge, Downing Place, Cambridge CB2 3EN, UK. ²⁹Centre for Past Climate Change and Department of Meteorology, University of Reading, Earley Gate, Post Office Box 243, Reading, UK. ³⁰School of Geographical Sciences, University Road, Clifton, Bristol BS8 1SS, UK. ³¹Escuela Nacional de Antropología e Historia, Periférico Sur y Zapote s/n Colonia Isidro Fabela, Tlalpan, Isidro Fabela, 14030 Mexico City, Mexico. ³²Instituto de Investigación en Genética Molecular, Universidad de Guadalajara, Ocotlán, Mexico. ³³Dipartimento di Scienze Biologiche, Geologiche e Ambientali (BiGeA), Università di Bologna, Via Selmi 3, 40126 Bologna, Italy. ³⁴Department of Anthropology, Oregon State University, 238 Waldo Hall, Corvallis, OR 97331 USA. ³⁵Museum National d'Histoire Naturelle, CNRS, Université Paris 7 Diderot, Sorbonne Paris Cité, Sorbonne Universités, Unité Eco-Anthropologie et Ethnobiologie (UMR7206), Paris, France. ³⁶Anthropological Institute and Museum, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland. ³⁷Department of Archaeology, Memorial University, Queen's College, 210 Prince Philip Drive, St. John's, Newfoundland A1C 5S7, Canada. ³⁸Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, Leipzig 04103, Germany. ³⁹Department of Earth System Science, University of California, Irvine, Keck Carbon Cycle Accelerator Mass Spectrometry Group, B321 Croul Hall, Irvine, CA 92697, USA. ⁴⁰Instituto de Alta Investigación, Universidad de Tarapacá, 18 de Septiembre 2222, Carisilla 6-D Arica, Chile. ⁴¹Departamento de Antropología, Universidad de Tarapacá, 18 de Septiembre 2222, Carisilla 6-D Arica, Chile. ⁴²Fundação Museu do Homem Americano, Centro Cultural Sérgio Motta, Campeste, 64770-000 São Raimundo Nonato, Brazil. ⁴³Laboratoire d'Anthropologie Moléculaire et Imagerie de Synthèse UMR-5288, CNRS, Université de Toulouse, 31073 Toulouse, France. ⁴⁴Institute of Cytology and Genetics, Siberian Branch of the Russian Academy of Sciences, Prospekt Lavrentyeva 10, 630090 Novosibirsk, Russia. ⁴⁵Novosibirsk State University, 2 Pirogova Street, 630090 Novosibirsk, Russia. ⁴⁶Institute of Internal Medicine, Siberian Branch of RAS, 175/1 ul. B. Bogatkova, Novosibirsk 630089, Russia. ⁴⁷Novosibirsk State University, Laboratory of Molecular Epidemiology and Bioinformatics, 630090 Novosibirsk, Russia. ⁴⁸Vavilov Institute of General Genetics, Gubkina 3, 119333 Moscow, Russia. ⁴⁹Research Centre for Medical Genetics, Moskvorechie 1, 115478 Moscow, Russia. ⁵⁰Kemerovo State University, Krasnaya 3, 650000 Kemerovo, Russia. ⁵¹Institute of Biochemistry and Genetics, Ufa Scientific Center of Russian Academy of Sciences, Prospekt Oktyabrya 71, 450054 Ufa, Russia. ⁵²Department of Genetics and Fundamental Medicine, Bashkir State University, Zaki Validi 32, 450076 Ufa, Russia. ⁵³Department of Molecular Genetics, Yakut Scientific Centre of Complex Medical Problems, Sergelyahskoe Shosse 4, 677010 Yakutsk, Russia. ⁵⁴Laboratory of Molecular Biology, Institute of Natural Sciences, M. K. Ammosov North-Eastern Federal University, 677000 Yakutsk, Russia. ⁵⁵Institute of Biological Problems of the North, Russian Academy of Sciences, Portovaya Street 18, Magadan 685000, Russia. ⁵⁶Department of Anthropology, Western Washington University, Bellingham, WA 98225, USA. ⁵⁷Department of Anthropology, Northwest Community College, 353 Fifth Street, Prince Rupert, British Columbia V8J 3L6, Canada. ⁵⁸Canadian Museum of History, 100 Rue Laurier, Gatineau, Quebec K1A 0M8, Canada. ⁵⁹University of Western Ontario, London, Ontario N6A 3K7, Canada. ⁶⁰Simon Fraser University, Burnaby, British Columbia V5A 1S6, Canada. ⁶¹Metlakatla Treaty Office, Post Office Box 224, Prince Rupert, BC V8J 3P6, Canada. ⁶²Sealaska Heritage Institute, 105 S. Seward Street, Juneau, AK 99801, USA. ⁶³Department of Structural Biology, Stanford University School of Medicine, D100 Fairchild Science Building, Stanford, CA 94305-5126, USA. ⁶⁴Department of Anthropology, Washington State University, Pullman, WA 99163, USA. ⁶⁵Division of Biological Anthropology, University of Cambridge, Henry Wellcome Building, Fitzwilliam Street, Cambridge CB2 1QH, UK. ⁶⁶Department of Medicine, University of Cambridge, Medical Research Council Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge Biomedical Campus, Cambridge CB2 0QH, UK. ⁶⁷Laboratory of Biological Anthropology, University of Kansas, 1415 Jayhawk Boulevard, 622 Fraser Hall, Lawrence, KS 66045, USA. ⁶⁸Molecular Anthropology Laboratory, 209 Young Hall, Department of Anthropology, University of California, One Shields Avenue, Davis, CA 95616, USA. ⁶⁹Center for the Study of the First Americans, Texas A&M University, College Station, TX 77843-4352, USA. ⁷⁰Department of Anthropology, Texas A&M University, College Station, TX 77843-4352, USA. ⁷¹Department of Geography, Texas A&M University, College Station, TX 77843-4352, USA. ⁷²Santa Barbara Museum of Natural History, 2559 Puesta del Sol, Santa Barbara, CA 93105, USA. ⁷³Carle R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA. ⁷⁴Department of Anthropology, Southern Methodist University, Dallas, TX 75275, USA. ⁷⁵Department of Integrative Biology, University of California, 3060 Valley Life Sciences Building 3140, Berkeley, CA 94720, USA.

*These authors contributed equally to this work. †Corresponding authors. E-mail: ewillierslev@snm.ku.dk (E.W.); rasmus_nielsen@berkeley.edu (R.N.); jss@berkeley.edu (Y.S.S.)




Fig. 1. Origins and population history of Native Americans. (A) Our results show that the ancestors of all present-day Native Americans, including Amerindians and Athabascans, derived from a single migration wave into the Americas (purple), separate from the Inuit (green). This migration from East Asia occurred no earlier than 23 ka and is in agreement with archaeological evidence from sites such as Monte Verde (50). A split between the northern and southern branches of Native Americans occurred ~13 ka, with the former comprising Athabascans and northern Amerindians and the latter consisting of Amerindians in southern North America and Central and South America, including the Anzick-1 individual (5). There is an admixture signal between Inuit and Athabascans as well as some northern Amerindians (yellow line); however, the gene flow direction is unresolved because of the complexity of the admixture events (28). Additionally, we see a weak signal related to Australo-Melanesians in some Native Americans, which may have been mediated through East Asians and Aleutian Islanders (yellow arrows). Also shown is the Mal'ta gene flow into Native American ancestors some 23 ka (yellow arrow) (4). It is currently not possible for us to ascertain the exact geographical locations of the depicted events; hence, the positioning of the arrows should not be considered a reflection of these. (B) Admixture plot created on the basis of TreeMix results (fig. S5) shows that all Native Americans form a clade, separate from the Inuit, with gene flow between some Native Americans and the North American Arctic. The number of genome-sequenced individuals included in the analysis is shown in brackets.

resolved because of the complexity of the admixture events (28). Additionally, we see a weak signal related to Australo-Melanesians in some Native Americans, which may have been mediated through East Asians and Aleutian Islanders (yellow arrows). Also shown is the Mal'ta gene flow into Native American ancestors some 23 ka (yellow arrow) (4). It is currently not possible for us to ascertain the exact geographical locations of the depicted events; hence, the positioning of the arrows should not be considered a reflection of these. (B) Admixture plot created on the basis of TreeMix results (fig. S5) shows that all Native Americans form a clade, separate from the Inuit, with gene flow between some Native Americans and the North American Arctic. The number of genome-sequenced individuals included in the analysis is shown in brackets.

first estimated divergence times between Native Americans and the Koryak of Siberia, one of the genetically closest sampled East Asian populations to Native Americans (fig. S5), using demographic models that reflect a clean split between the populations (28). With both diCal2.0 and the IBS tract method, the split of Native Americans (including Amerindians and Athabascans) from the Koryak dates to ~20 ka (tables S11A and S12 and fig. S15) (28).

We further applied diCal2.0 to models with gene flow postdating the split between Native Americans and Koryak (Fig. 2A) and found that they provided a better fit to the data than did the models without gene flow (fig. S18) (28). Overall, simulated data based on the models inferred by using diCal2.0 and real data show very similar IBS tract length distributions (Fig. 2B) and relative cross coalescence rates (CCRs) between pairs of individuals estimated by using the Multiple Se-

quentially Markovian Coalescent (MSMC) method (method 3) (Fig. 2, C and D) (28, 44). This serves as a confirmation for the model estimates from diCal2.0. We evaluated all three methods using simulations under complex demographic models and additionally investigated the effects of switch-errors in haplotype phasing on the estimates (28).

We then applied the diCal2.0 model that allows for gene flow between populations after




Fig. 2. Divergence estimates between Native Americans and Siberian Koryak. (A) The demographic model used allows for continuous gene flow between populations 1 and 2, starting from the time T_{DIV} of divergence and ending at T_M . The backward probability of migration per individual per generation is denoted by m . The bottleneck at T_B captures the out-of-Africa event. (B) The red and black solid curves depict empirical distributions of IBS tracts shared between Karitiana-Koryak and Athabascan-Koryak, respectively. The orange, pink, dashed blue, and dashed green curves depict IBS tracts shared between the two population pairs, simulated under two demographic models based on results from diCal2.0. Overall, for Karitiana-Koryak and Athabascan-

Koryak, the migration scenarios (orange and pink, respectively) match the empirical curves (red and black, respectively) better than the clean split scenarios match (dashed blue and dashed green, respectively), with more long IBS tracts showing evidence of recent common ancestry between Koryaks and Native Americans. (C and D) Relative CCRs for the Karitiana-Koryak and Athabascan-Koryak divergence (red), respectively, including data simulated under the two demographic models in (B). In both cases, the model with gene flow (orange) fits the data (red) better than does the clean split model (blue). The migration model explains a broader CCR tail in the case of Karitiana-Koryak and the relatively late onset of the CCR decay for Athabascan-Koryak.

their split in order to estimate divergence times for Native Americans from more geographically and genetically distant East Asian groups, including the Siberian Nivkh and Han Chinese. As before, the divergence estimates for Amerindians and Athabascans were very similar to one another, ~23 ka (table S11B and figs. S18 and S21).

Hence, our results suggest that Amerindians and Athabascans were, by three different methods, consistently equidistant in time to populations that were sampled from different regions of East Asia, including some proximate to Beringia, and with varied population histories. This suggests that these two major Native American subgroups are descendants of the same source

population that split off from ancestral East Asians during the LGM. It is conceivable that harsh climatic conditions during the LGM may have contributed to the isolation of ancestral Native Americans, ultimately leading to their genetic divergence from their East Asian ancestors.

We also modeled the peopling of the Americas using a climate-informed spatial genetic model (CISGeM), in which the genetic history and local demography is informed by paleoclimatic and paleovegetation reconstructions (28, 45), and found the results to be in accordance with the conclusion of a single migration source for all Native Americans. Using present-day and ancient high-coverage genomes, we found that

Athabascans and Anzick-1, but not Greenlandic Inuit and Saqqaq (29, 39), belong to the same initial migration wave that also gave rise to present-day Amerindians from southern North America and Central and South America (Fig. 3) and that this migration likely followed a coastal route, given our current understanding of the glacial geological and paleoenvironmental parameters of the Late Pleistocene (fig. S31).

In all cases, the best fit of the demographic models to the IBS tract distribution and relative CCR by MSMC required gene flow between Siberian and Native American populations after their initial split (Fig. 2, B to D). We also found strong evidence for gene flow between Athabascans and




Fig. 3. Testing migrations into the Americas by using a climate-informed model. Estimates of difference in genetic divergence between Amerindians (from southern North America and Central and South America) or Koryak versus Athabaskan and Greenlandic Inuit and the ancient Saqqaq and Anzick-1 genomes (black vertical lines), compared with posterior probability distribution predicted from a climate-informed spatial genetic model reconstructing a single wave into

the Americas (curves, the colored part represents the 95% credibility interval). ΔT for population X is defined as $T(X, \text{Koryak}) - T(X, \text{Central and South Amerindians})$ (28). Both Anzick-1 and the Athabascans were part of the same wave into the Americas to which other Amerindian populations from southern North America and Central and South America belonged, whereas the Inuit and Saqqaq are the descendants of different waves (observed values outside the 95% credibility interval).

the Inuit (table S11B), supported by results from ADMIXTURE (fig. S4), TreeMix (fig. S5), D -statistics using both whole-genome and SNP chip genotype data (figs. S6 and S8A) (28, 46, 47), and outgroup f_3 -statistics using whole-genome data (fig. S12) (28, 47). We attempted to estimate the divergence times between Inuit and Siberians as well as Inuit and Native Americans (table S11 and figs. S19 and S25 to S27), but our analyses were complicated by gene flow between Inuit and Athabascans as well as complex admixture patterns among Arctic groups (fig. S5).

We tested the duration and magnitude of post-split gene flow between Native Americans and Siberians using diCal2.0 by introducing stopping time of gene flow as a free parameter (28). We still obtained the highest likelihood for a divergence time of 22 ka between Amerindians and Siberians as well as Athabascans and Siberians, although estimates for gene-flow rate and end of the gene flow differ (table S11C and fig. S22). Gene flow between Athabascans and Siberians seems to have stopped ~12 ka (table S11C), suggesting a link to the breaching of the Beringian Land Bridge by rising sea levels (48).

Overall, our results support a common Siberian origin for all Native Americans, contradicting claims for an early migration to the Americas from Europe (49), with their initial isolation and entrance into the Americas occurring no earlier than 23 ka, but with subsequent admixture with East Asian populations. This additionally suggests that the Mal'ta-related admixture into the early Americans (4), representing ancestors of both Amerindians and Athabascans (Fig. 1 and fig. S5), occurred sometime after 23 ka, after the Native American split from East Asians.

Subsequent in situ diversification of Native American groups

That Amerindian and Athabaskan groups were part of the same migration implies that present-day genetic differences observed between them must have arisen later, after ~23 ka. Using the clean-split model in diCal2.0 on the modern genomes data set, we estimated that Athabascans

and Karitiana diverged ~13 ka (95% confidence interval of ~11.5 to 14.5 ka, estimated from parametric bootstrap results) (table S11A and fig. S16), which is consistent with results from MSMC (fig. S27) (28).

Where the divergence between Karitiana and Athabascans occurred is not known. However, several independent lines of evidence suggest that it is more likely to have occurred in lower-latitude North America instead of eastern Beringia (Alaska). These include the equidistant split times of Amerindians and Athabascans to Asian populations, the relatively brief interval between their estimated divergence date range and the age of Anzick-1 (12.6 ka) (5), and last, the geographic location of Anzick-1 to the south of the North American ice sheets and its clear affiliation with the “southern branch” of Native Americans (taken broadly to include Amerindians from southern North America and Central and South America) (5), as determined with outgroup f_3 -statistics by using SNP chip genotype data from present-day worldwide populations (Fig. 4 and figs. S13 and S14) (47). Divergence in North America would also be consistent with the known pre-Clovis age sites in the Americas, such as Monte Verde (14.6 ka) (50). The most parsimonious model would be that both Amerindians and Athabascans are descendants of the same ancestral Native American population that entered the Americas then subsequently diversified. However, we cannot discount alternative and more complex scenarios, which could be tested with additional ancient samples.

By the Clovis period (~12.6 ka), the ancestral Native American population had already diversified into “northern” and “southern” branches, with the former including ancestors of present-day Athabascans and northern Amerindian groups such as Chipewyan, Cree, and Ojibwa and the latter including Amerindians from southern North America and Central and South America (Fig. 4 and fig. S14). We tested whether later gene flow from East Asian sources, such as the Inuit, might explain the genetic differences between these two branches. Using D -statistics

on SNP chip genotype data (47) masked for non-native ancestry, we observed a signal of gene flow between the Inuit and northwest Pacific Coast Amerindians such as Coastal Tsimshian and Nisga'a, residing in the same region as the northern Athabascans (fig. S8B) (28). However, this signal of admixture with the Inuit, also detected in Athabascans (figs. S6 and S8A), was not evident among northern Amerindian populations located further east, such as Cree, Ojibwa, and Chipewyan (fig. S8C) (28). This suggests that the observed difference between the northern and southern branches is not a consequence of post-split East Asian gene flow into the northern branch and also provides a possible explanation as to why the southern branch Amerindians such as Karitiana are genetically closer to the northern Amerindians located further east than to northwest coast Amerindians and Athabascans (fig. S9).

In contrast to Anzick-1, several of the Holocene individuals from the Americas—including those sequenced in this study, as well as the 8500-year-old Kennewick Man (51)—are closely related to present-day Native American populations from the same geographical regions (Fig. 4 and figs. S13 and S14). This implies genetic continuity of ancient and modern populations in some parts of the Americas over at least the past 8500 years, which is in agreement with recent results from Kennewick Man (51).

Evidence of more distant Old World gene flow into some Native Americans

When testing for gene flow between Athabascans and Inuit with masked SNP chip genotype data-based D -statistics (47) (fig. S8), we observed a weak tendency for the Inuit to be much closer to the Athabascans than to certain Amerindians such as the North American Algonquin and Cree, and the Yaqui and Arhuaco of Central and South America (respectively), as compared with other Amerindians such as the Palikur and Surui of Brazil (fig. S8).

To further investigate this trend, we tested for additional gene flow from Eurasian populations into the Americas with D -statistics using the




Fig. 4. Diversification within the Americas. SNP chip genotype data-based outgroup f_3 -statistics (47) of the form $f_3(X, \text{Ancient}; \text{Yoruba})$ were used to estimate the shared ancestry between ancient samples from the Americas and a large panel of worldwide present-day populations (X), including Athabascan and Amerindian groups from North America (table S3), some of which were masked for non-native ancestry before the analysis (28). The outgroup f_3 -statistics are depicted as heat maps, with the sampling location of the ancient sample

marked by the dotted lines, and corresponding ranked plots with error bars are shown in fig. S14. “BP” refers to time before present. We find the Anzick-1 sample to share most ancestry with the southern branch of Native Americans when using multiple northern Native Americans sequenced in this study, which is consistent with (5). The seven Holocene age samples share most ancestry with Native Americans, with a general tendency to be genetically closer to present-day Native American populations from the same geographical region.

masked SNP chip genotype data set (47). We found that some American populations—including the Aleutian Islanders, Surui, and Athabascans—are closer to Australo-Melanesians as compared with other Native Americans, such as North American Ojibwa, Cree, and Algonquin and the South American Purepecha, Arhuaco, and Wayuu (fig. S10). The Surui are, in fact, one of closest Native American populations to East Asians and Australo-Melanesians, the latter including Papuans, non-Papuan Melanesians, Solomon Islanders, and South East Asian hunter-gatherers such as Aeta (fig. S10). We acknowledge that this observation is based on the analysis of a small fraction of the whole-genome and SNP chip genotype data sets—especially for the Aleutian Islander data, which is heavily masked owing to recent admixture with Europeans (28)—and that the trends in the data are weak.

Nonetheless, if it proves correct, these results suggest that there may be a distant Old World signal related to Australo-Melanesians and East Asians in some Native Americans. The widely scattered and differential affinity of Native Americans to the Australo-Melanesians, ranging from a strong signal in the Surui to a much weaker signal in northern Amerindians such as Ojibwa, points to this gene flow occurring after the initial peopling by Native American ancestors.

However, how this signal may have ultimately reached South America remains unclear. One possible means is along a northern route via the Aleutian Islanders, previously found to be closely related to the Inuit (39), who have a relatively greater affinity to East Asians, Oceanians, and Denisovan than Native Americans in both whole-genome and SNP chip genotype data-based *D* tests (table S10 and figs. S10 and S11). On the basis of archaeological evidence and mtDNA data from ancient and modern samples, the Aleutian Islands are hypothesized to have been peopled as early as ~9 ka by “Paleo-Aleuts” who were succeeded by the “Neo-Aleuts,” with present-day Aleutian Islanders potentially resulting from admixture between these two populations (52, 53). Perhaps their complex genetic history included input from a population related to Australo-Melanesians through an East Asian continental route, and this genomic signal might have been subsequently transferred to parts of the Americas, including South America, through past gene flow events (Fig. 1). Evidence for this gene flow is supported with diCal2.0 and MSMC analyses showing a weak but recent gene flow into South Americans from populations related to present-day Northeast Asians (Koryak) (Fig. 2C and table SIIC), who might be considered a proxy for the related Aleutian Islanders.

Testing the Paleoamerican model

The detection of an Australo-Melanesian genetic signal in the Americas, however subtle, returns the discussion to the Paleoamerican model, which hypothesizes, on the basis of cranial morphology, that two temporally and source-distinct populations colonized the Americas. The earlier population reportedly originated in Asia in the Late

Pleistocene and gave rise to both Paleoamericans and present-day Australo-Melanesians, whose shared cranial morphological attributes are presumed to indicate their common ancestry (23). The Paleoamericans were, in turn, thought to have been largely replaced by ancestors of present-day Amerindians, whose crania resemble modern East Asians and who are argued to be descendants of later arriving Mongoloid populations (14, 23, 26, 54). The presence of Paleoamericans is inferred primarily from ancient archaeological specimens in North and South America and a few relict populations of more recent age, which include the extinct Pericúes and Fuego-Patagonians (24, 25, 55).

The Paleoamerican hypothesis predicts that these groups should be genetically closer to Australo-Melanesians than other Amerindians. Previous studies of mtDNA and Y chromosome data obtained from Fuego-Patagonian and Paleoamerican skeletons have identified haplogroups similar to those of modern Native Americans (55–57). Although these results indicate some shared maternal and paternal ancestry with contemporary Native Americans, uniparental markers can be misleading when drawing conclusions about the demographic history of populations. To conclusively identify the broader population of ancestors who may have contributed to the Paleoamerican gene pool, autosomal genomic data are required.

We therefore sequenced 17 ancient individuals affiliated to the now-extinct Pericúes from Mexico and Fuego-Patagonians from Chile and Argentina (28), who, on the basis of their distinctive skull morphologies, are claimed to be relicts of Paleoamericans (23, 27, 58, 59). Additionally, we sequenced two pre-Columbian mummies from northern Mexico (Sierra Tarahumara) to serve as morphological controls because they are expected to fall within the range of Native American morphological cranial variation (28). We found that the ancient samples cluster with other Native American groups and are outside the range of Oceanian genetic variation (Fig. 5 and figs. S32, S33, and S34) (28). Similarly, outgroup f_3 -statistics (47) reveal low shared genetic ancestry between the ancient samples and Oceanians (figs. S36 and S37) (28), and genome-based and masked SNP chip genotype data-based *D*-statistics (46, 47) show no evidence for gene flow from Oceanians into the Pericúes or Fuego-Patagonians (fig. S39) (28).

Because the Paleoamerican model is based on cranial morphology (23, 27, 58, 59), we also measured craniometric data for the ancient samples and assessed their phenotypic affinities to supposed Paleoamericans, Amerindians, and worldwide populations (28). The results revealed that the analyzed Fuego-Patagonians showed closest craniometric affinity to Arctic populations and the Paleoamericans, whereas the analyzed female Pericúes showed closest craniometric affinities to populations from North America, the Arctic region, and Northern Japan (table SI5). Our analyses demonstrated that the presumed ancestral ancient Paleoamerican reference sample from Lagoa Santa, Brazil (24) had closest affinities to Arctic and East Asian populations (table SI5).

Consequently, for the Fuego-Patagonians, the female Pericúes, and the Lagoa Santa Paleoamerican sample, we were not able to replicate previous results (24) that report close similarity of Paleoamerican and Australo-Melanesian cranial morphologies. Male Pericúes samples displayed more craniometric affinities with populations from Africa and Australia relative to the female individuals of their population (fig. S41). The results of analyses based on craniometric data thus are highly sensitive to sample structure and the statistical approach and data filtering used (51). Our morphometric analyses suggest that these ancient samples are not true relicts of a distinct migration as claimed and hence do not support the Paleoamerican model. Similarly, our genomic data also provide no support for an early migration of populations directly related to Australo-Melanesians into the Americas.

Discussion

That Native Americans diverged from their East Asian ancestors during the LGM and no earlier than 23 ka provides an upper bound, and perhaps the climatic and environmental context, for the initial isolation of their ancestral population and a maximum estimate for the entrance and subsequent spread into the Americas. This result is consistent with the model that people entered the Americas before the development of the Clovis complex and had reached as far as southern South America by 14.6 ka. Because archaeological evidence provides only a minimum age for human presence in the Americas, we can anticipate the possible discovery of sites that approach the time of the divergence of East Asians and Native Americans. However, our estimate for the initial divergence and entry of Native American ancestors does not support archaeological claims for an initial peopling substantially earlier than the LGM (8–10).

Although our data cannot provide the precise geographical context for the initial peopling process, it has allowed us to more accurately estimate its temporal dynamics. This, in turn, has enabled us to reassess the Beringian Incubation Model, which, based on mtDNA data and the timing and geographical distribution of archaeological sites, hypothesized a ~15,000-year-long period of isolation of ancestral Native Americans in Beringia during the LGM (19–21). Our results, along with recent findings of mtDNA haplogroup C1 in Iceland and ancient northwest Russia (60), do not fit with the proposed 15,000-year span of the Beringian Incubation Model (19–21). It is possible that a shorter period of isolation occurred (~8000 years), but whether it occurred in Siberia or Beringia will have to be determined from future ancient DNA and archaeological findings. Given the genetic continuity between Native Americans and some East Asian populations (figs. S4 and S5), other demographic factors, such as surfing during population expansions into unoccupied regions (61), may ultimately need to be taken into account to better understand the presence of a large number of high-frequency private variants in the indigenous populations of the Americas.




Fig. 5. The Paleoamerican model. (A) Principal component analysis plot of 19 ancient samples combined with a worldwide reference panel, including 1823 individuals from (6). Our samples plot exclusively with American samples. Plots with other reference panels consisting of Native American populations are provided in fig. S32. (B) Population structure in the ancient Pericú, Mexican mummy, and Fuego-Patagonian individuals from this study. Ancestry proportions

are shown when assuming six ancestral populations ($K = 6$). The top bar shows the ancestry proportions of the 19 ancient individuals, Anzick-1 (5), and two present-day Native American genomes from this study (Huichol and Aymara). The plot at the bottom illustrates the ancestry proportions for 1823 individuals from (6). Our samples show primarily Native American (ivory, >92%) and Siberian (red, ~5%) ancestry. The plot with $K = 13$ is provided in fig. S33.

The data presented here are consistent with a single initial migration of all Native Americans and with later gene flow from sources related to East Asians and, more distantly, Australo-

Melanesians. From that single migration, there was a diversification of ancestral Native Americans leading to the formation of northern and southern branches, which appears to have taken

place ~13 ka within the Americas. This split is consistent with the patterns of uniparental genomic regions of mtDNA haplogroup X and some Y chromosome C haplotypes being present in

northern, but not southern, populations in the Americas (18, 62). This diversification event coincides roughly with the opening of habitable routes along the coastal and the interior corridors into unglaciated North America some 16 and 14 ka, respectively (63, 64), suggesting a possible role of one or both of these routes in the isolation and subsequent dispersal of Native Americans across the continent.

Methods

DNA was extracted from 31 present-day individuals from the Americas, Siberia, and Oceania and 23 ancient samples from the Americas and converted to Illumina libraries and shotgun sequenced (28). Three of the ancient samples were radiocarbon dated, of which two were corrected for marine reservoir offset (28). SNP chip genotype data was generated from 79 present-day Siberians and Native Americans affiliated to 28 populations (28). Raw data from SNP chip and shotgun sequencing were processed by using standard computational procedures (28). Error rate analysis, DNA damage analysis, contamination estimation, sex determination, mtDNA and Y chromosome haplogroup assignment, ADMIXTURE analysis, ancestry painting and admixture masking, principal component analysis using SNP chip genotype data, TreeMix analysis on genomic sequence data, D -statistic and outgroup f_3 -statistic tests on SNP chip genotype and genomic sequence data, divergence time estimation by use of diCal2.0, an IBS tract method and MSMC, climate-informed spatial genetic model analysis, and craniometric analysis were performed as described (28).

REFERENCES AND NOTES

1. T. D. Dillehay, The late Pleistocene cultures of South America. *Evol. Anthropol.* **7**, 206–216 (1999). doi: [10.1002/\(SICI\)1520-6505\(1999\)7:6<206::AID-EVAN5>3.0.CO;2-G](#)
2. D. L. Jenkins et al., Clovis age Western Stemmed projectile points and human coprolites at the Paisley Caves. *Science* **337**, 223–228 (2012). PMID: [22798611](#)
3. D. J. Meltzer, *First Peoples in a New World: Colonizing Ice Age America* (Univ. California Press, Berkeley, 2009).
4. M. Raghavan et al., Upper Palaeolithic Siberian genome reveals dual ancestry of Native Americans. *Nature* **505**, 87–91 (2014). doi: [10.1038/nature12736](#); PMID: [24256729](#)
5. M. Rasmussen et al., The genome of a Late Pleistocene human from a Clovis burial site in western Montana. *Nature* **506**, 225–229 (2014). doi: [10.1038/nature13025](#); PMID: [24522598](#)
6. D. Reich et al., Reconstructing Native American population history. *Nature* **488**, 370–374 (2012). PMID: [22801491](#)
7. M. R. Waters et al., The Buttermilk Creek complex and the origins of Clovis at the Debra L. Friedkin site, Texas. *Science* **331**, 1599–1603 (2011). doi: [10.1126/science.1201855](#); PMID: [21436451](#)
8. G. M. Santos et al., A revised chronology of the lowest occupation layer of Pedra Furada Rock Shelter, Piauí, Brazil: The Pleistocene peopling of the Americas. *Quat. Sci. Rev.* **22**, 2303–2310 (2003). doi: [10.1016/S0277-3791\(03\)00205-1](#)
9. S. R. Holen, K. Holen, K. in *Paleoamerican Odyssey*, K. E. Graf, C. V. Ketron, M. R. Waters, Eds. (Texas A&M Univ. Press, College Station, 2014), pp. 429–444.
10. E. Boëda et al., A new late Pleistocene archaeological sequence in South America: The Vale da Pedra Furada (Piauí, Brazil). *Antiquity* **88**, 927–941 (2014). doi: [10.1017/S0003598X00050845](#)
11. D. W. Owsley, R. L. Jantz, *Kennewick Man: The Scientific Investigation of an Ancient American Skeleton* (Texas A&M Univ. Press, College Station, 2014).
12. A. Achilli et al., The phylogeny of the four pan-American mtDNA haplogroups: Implications for evolutionary and disease studies. *PLOS ONE* **3**, e1764 (2008). doi: [10.1371/journal.pone.0001764](#); PMID: [18335039](#)
13. V. Battaglia et al., The first peopling of South America: New evidence from Y-chromosome haplogroup Q. *PLOS ONE* **8**, e71390 (2013). doi: [10.1371/journal.pone.0071390](#); PMID: [23990949](#)
14. C. L. Brace et al., Old World sources of the first New World human inhabitants: A comparative craniofacial view. *Proc. Natl. Acad. Sci. U.S.A.* **98**, 10017–10022 (2001). doi: [10.1073/pnas.171305898](#); PMID: [11481450](#)
15. U. A. Perego et al., Distinctive Paleo-Indian migration routes from Beringia marked by two rare mtDNA haplogroups. *Curr. Biol.* **19**, 1–8 (2009). doi: [10.1016/j.cub.2008.11.058](#); PMID: [19135370](#)
16. U. A. Perego et al., The initial peopling of the Americas: A growing number of founding mitochondrial genomes from Beringia. *Genome Res.* **20**, 1174–1179 (2010). doi: [10.1101/gr.109231.110](#); PMID: [20587512](#)
17. N. J. R. Fagundes, R. Kanitz, S. L. Bonatto, A reevaluation of the Native American mtDNA genome diversity and its bearing on the models of early colonization of Beringia. *PLOS ONE* **3**, e3157 (2008). doi: [10.1371/journal.pone.0003157](#); PMID: [18797501](#)
18. S. L. Zegura, T. M. Karafet, L. A. Zhivotovskiy, M. F. Hammer, High-resolution SNPs and microsatellite haplotypes point to a single, recent entry of Native American Y chromosomes into the Americas. *Mol. Biol. Evol.* **21**, 164–175 (2004). doi: [10.1093/molbev/msh009](#); PMID: [14595095](#)
19. E. Tamm et al., Beringian standstill and spread of Native American founders. *PLOS ONE* **2**, e829 (2007). doi: [10.1371/journal.pone.0000829](#); PMID: [17786201](#)
20. A. Kitchen, M. M. Miyamoto, C. J. Mulligan, A three-stage colonization model for the peopling of the Americas. *PLOS ONE* **3**, e1596 (2008). doi: [10.1371/journal.pone.0001596](#); PMID: [18270583](#)
21. C. J. Mulligan, A. Kitchen, M. M. Miyamoto, Updated three-stage model for the peopling of the Americas. *PLOS ONE* **3**, e3199 (2008). doi: [10.1371/journal.pone.0003199](#); PMID: [18797500](#)
22. K. B. Schroeder et al., A private allele ubiquitous in the Americas. *Biol. Lett.* **3**, 218–223 (2007). doi: [10.1098/rsbl.2006.0609](#); PMID: [17301009](#)
23. R. González-José et al., Craniometric evidence for Palaeoamerican survival in Baja California. *Nature* **425**, 62–65 (2003). doi: [10.1038/nature01816](#); PMID: [12955139](#)
24. W. A. Neves, M. Hubbe, Cranial morphology of early Americans from Lagoa Santa, Brazil: Implications for the settlement of the New World. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 18309–18314 (2005). doi: [10.1073/pnas.0507185102](#); PMID: [16344464](#)
25. W. Neves et al., in *Paleoamerican Odyssey*, K. E. Graf, C. V. Ketron, M. R. Waters, Eds. (Texas A&M Univ. Press, College Station, 2014), pp. 397–412.
26. R. González-José, M. C. Bortolini, F. R. Santos, S. L. Bonatto, The peopling of America: Craniofacial shape variation on a continental scale and its interpretation from an interdisciplinary view. *Am. J. Phys. Anthropol.* **137**, 175–187 (2008). doi: [10.1002/ajpa.20854](#); PMID: [18481303](#)
27. M. Lahr, Patterns of modern human diversification: Implications for Amerindian origins. *Am. J. Phys. Anthropol.* **38** (S21), 163–198 (1995). doi: [10.1002/ajpa.1330380609](#)
28. Materials and methods are available as supplementary materials on Science Online.
29. M. Rasmussen et al., Ancient human genome sequence of an extinct Palaeo-Eskimo. *Nature* **463**, 757–762 (2010). doi: [10.1038/nature08835](#); PMID: [20148029](#)
30. B. Yunusbayev et al., The genetic legacy of the expansion of Turkic-speaking nomads across Eurasia. *PLOS Genet.* **11**, e1005068 (2015). doi: [10.1371/journal.pgen.1005068](#); PMID: [25989006](#)
31. A. Cardona et al., Genome-wide analysis of cold adaptation in indigenous Siberian populations. *PLOS ONE* **9**, e98076 (2014). PMID: [24847810](#)
32. J. Z. Li et al., Worldwide human relationships inferred from genome-wide patterns of variation. *Science* **319**, 1100–1104 (2008). doi: [10.1126/science.1153717](#); PMID: [18292342](#)
33. A. Moreno-Estrada et al., Reconstructing the population genetic history of the Caribbean. *PLOS Genet.* **9**, e1003925 (2013). doi: [10.1371/journal.pgen.1003925](#); PMID: [24244192](#)
34. A. Moreno-Estrada et al., The genetics of Mexico recapitulates Native American substructure and affects biomedical traits. *Science* **344**, 1280–1285 (2014). doi: [10.1126/science.1251688](#); PMID: [24926019](#)
35. P. Verdu et al., Patterns of admixture and population structure in native populations of Northwest North America. *PLOS Genet.* **10**, e1004530 (2014). doi: [10.1371/journal.pgen.1004530](#); PMID: [25122539](#)
36. D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009). doi: [10.1101/gr.094052.109](#); PMID: [19648217](#)
37. J. K. Pickrell, J. K. Pritchard, Inference of population splits and mixtures from genome-wide allele frequency data. *PLOS Genet.* **8**, e1002967 (2012). doi: [10.1371/journal.pgen.1002967](#); PMID: [23166502](#)
38. M. Meyer et al., A high-coverage genome sequence from an archaic Denisovan individual. *Science* **338**, 222–226 (2012). PMID: [22936568](#)
39. M. Raghavan et al., The genetic prehistory of the New World Arctic. *Science* **345**, 1255832–1255832 (2014). doi: [10.1126/science.1255832](#); PMID: [25170159](#)
40. A. Scally, R. Durbin, Revising the human mutation rate: Implications for understanding human evolution. *Nat. Rev. Genet.* **13**, 745–753 (2012). doi: [10.1038/nrg3295](#); PMID: [22695354](#)
41. S. Sheehan, K. Harris, Y. S. Song, Estimating variable effective population sizes from multiple genomes: A sequentially markov conditional sampling distribution approach. *Genetics* **194**, 647–662 (2013). doi: [10.1534/genetics.112.149096](#); PMID: [23608192](#)
42. M. Steinrücken, J. S. Paul, Y. S. Song, A sequentially Markov conditional sampling distribution for structured populations with migration and recombination. *Theor. Popul. Biol.* **87**, 51–61 (2013). doi: [10.1016/j.tpb.2012.08.004](#); PMID: [23010245](#)
43. K. Harris, R. Nielsen, Inferring demographic history from a spectrum of shared haplotype lengths. *PLOS Genet.* **9**, e1003521 (2013). doi: [10.1371/journal.pgen.1003521](#); PMID: [23754952](#)
44. S. Schiffels, R. Durbin, Inferring human population size and separation history from multiple genome sequences. *Nat. Genet.* **46**, 919–925 (2014). doi: [10.1038/ng.3015](#); PMID: [24952747](#)
45. A. Eriksson et al., Late Pleistocene climate change and the global expansion of anatomically modern humans. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 16089–16094 (2012). doi: [10.1073/pnas.1209494109](#); PMID: [22988099](#)
46. R. E. Green et al., A draft sequence of the Neandertal genome. *Science* **328**, 710–722 (2010). doi: [10.1126/science.1188021](#); PMID: [20448178](#)
47. N. Patterson et al., Ancient admixture in human history. *Genetics* **192**, 1065–1093 (2012). doi: [10.1534/genetics.112.145037](#); PMID: [22960212](#)
48. J. F. Hoffecker, S. A. Elias, *Human Ecology of Beringia* (Columbia Univ. Press, New York, 2007).
49. S. Oppenheimer, B. Bradley, D. Stanford, Solutrean hypothesis: Genetics, the mammoth in the room. *World Archaeol.* **46**, 752–774 (2014). doi: [10.1080/00438243.2014.966273](#)
50. T. D. Dillehay, Monte Verde, A Late Pleistocene Settlement in Chile: The Archaeological Context and Interpretation (Smithsonian Institution Press, Washington DC, 1997).
51. M. Rasmussen et al., The ancestry and affiliations of Kennewick Man. *Nature* **10.1038/nature14625** (2015). doi: [10.1038/nature14625](#); PMID: [26087396](#)
52. R. S. Davis, R. A. Neech, Continuity and change in the eastern Aleutian archaeological sequence. *Hum. Biol.* **82**, 507–524 (2010). PMID: [21417882](#)
53. M. H. Crawford, R. C. Rubicz, M. Zloturo, Origins of Aleuts and the genetic structure of populations of the archipelago: Molecular and archaeological perspectives. *Hum. Biol.* **82**, 695–717 (2010). PMID: [21417890](#)
54. M. Hubbe, W. A. Neves, K. Harvati, Testing evolutionary and dispersion scenarios for the settlement of the new world. *PLOS ONE* **5**, e11105 (2010). PMID: [20559441](#)
55. J. C. Chatters et al., Late Pleistocene human skeleton and mtDNA link Paleoamericans and modern Native Americans. *Science* **344**, 750–754 (2014). doi: [10.1126/science.1252619](#); PMID: [24833392](#)
56. J. García-Bour et al., Early population differentiation in extinct aborigines from Tierra del Fuego-Patagonia: Ancient mtDNA sequences and Y-chromosome STR characterization. *Am. J. Phys. Anthropol.* **123**, 361–370 (2004). doi: [10.1002/ajpa.10337](#); PMID: [15022364](#)
57. S. I. Perez, V. Bernal, P. N. Gonzalez, M. Sardi, G. G. Politis, Discrepancy between cranial and DNA data of early Americans: Implications for American peopling. *PLOS ONE* **4**, e5746 (2009). doi: [10.1371/journal.pone.0005746](#); PMID: [19478947](#)
58. M. Hernández, C. L. Fox, C. García-Moro, Fuegian cranial morphology: The adaptation to a cold, harsh environment.

- Am. J. Phys. Anthropol.* **103**, 103–117 (1997). doi: [10.1002/\(SICI\)1096-8644\(199705\)103:1<103::AID-AJPA7>3.0.CO;2-X](https://doi.org/10.1002/(SICI)1096-8644(199705)103:1<103::AID-AJPA7>3.0.CO;2-X); pmid: [9185954](https://pubmed.ncbi.nlm.nih.gov/9185954/)
59. R. González-José, S. L. Dahinten, M. A. Luis, M. Hernández, H. M. Pucciarelli, Craniometric variation and the settlement of the Americas: Testing hypotheses by means of R-matrix and matrix correlation analyses. *Am. J. Phys. Anthropol.* **116**, 154–165 (2001). doi: [10.1002/ajpa.1108](https://doi.org/10.1002/ajpa.1108); pmid: [11590587](https://pubmed.ncbi.nlm.nih.gov/11590587/)
60. C. Der Sarkissian *et al.*, Mitochondrial genome sequencing in Mesolithic North East Europe Unearths a new sub-clone within the broadly distributed human haplogroup C1. *PLOS ONE* **9**, e87612 (2014). doi: [10.1371/journal.pone.0087612](https://doi.org/10.1371/journal.pone.0087612); pmid: [24503968](https://pubmed.ncbi.nlm.nih.gov/24503968/)
61. L. Excoffier, N. Ray, Surfing during population expansions promotes genetic revolutions and structuration. *Trends Ecol. Evol.* **23**, 347–351 (2008). doi: [10.1016/j.tree.2008.04.004](https://doi.org/10.1016/j.tree.2008.04.004); pmid: [18502536](https://pubmed.ncbi.nlm.nih.gov/18502536/)
62. A. Achilli *et al.*, Reconciling migration models to the Americas with the variation of North American native mitogenomes. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 14308–14313 (2013). pmid: [23940335](https://pubmed.ncbi.nlm.nih.gov/23940335/)
63. E. J. Dixon, Late Pleistocene colonization of North America from Northeast Asia: New insights from large-scale paleogeographic reconstructions. *Quat. Int.* **285**, 57–67 (2013). doi: [10.1016/j.quaint.2011.02.027](https://doi.org/10.1016/j.quaint.2011.02.027)
64. C. A. S. Mandryk, H. Josenhans, D. W. Fedje, R. W. Mathewes, Late Quaternary paleoenvironments of Northwestern North America: Implications for inland versus coastal migration routes. *Quat. Sci. Rev.* **20**, 301–314 (2001). doi: [10.1016/S0277-3791\(00\)00115-3](https://doi.org/10.1016/S0277-3791(00)00115-3)
- ACKNOWLEDGMENTS**
- We thank J. Valdés for providing craniometric measurements of the Pericúes at the National Museum of Anthropology in México; A. Monteverde from Centro Instituto Nacional de Antropología e Historia, Baja California Sur and V. Laborde at the Musée de l'Homme in Paris for providing documentation on Pericú and Fuego-Patagonian samples, respectively; T. Gilbert, M. McCoy, C. Sarkissian, M. Sikora, and L. Orlando for helpful discussions and input; D. Yao and C. Barbieri for helping with the collection of the Aymara population sample; B. Henn and J. Kidd for providing early access to the Mayan sequencing data; Canadian Museum of History; Metlakatla and Lax Kw'alaams First Nations; Listuguj Mi'gmaq Band Council; A. Pye of TERRA Facility, Core Research Equipment & Instrument Training (CREAIT) Network at Memorial University; the Danish National High-throughput DNA Sequencing Centre (Copenhagen) for help with sequencing; and Fondation Jean Dausset-Centre d'Etude du Polymorphisme Humain (CEPH) for providing DNA for the Human Genome Diversity Project (HGDP) samples that were genome-sequenced in this study. This study was supported by several funding bodies: Lundbeck Foundation and the Danish National Research Foundation (Centre for GeoGenetics members); Wellcome Trust grant 098051 (S.S., A.B., Y.X., C.T.-S., M.S.S., and R.D.); Marie Curie Intra-European Fellowship-FP7-People-PIEF-GA-2009-255503 and the Transforming Human Societies Research Focus Areas Fellowship from La Trobe University (C.V.); George Rosenkranz Prize for Health Care Research in Developing Countries and National Science Foundation award DMS-1201234 (M.C.A.A.); Swiss National Science Foundation (PBSKP3_143529) (A.-S.M.); Ministerio de Ciencia e Innovación (MICINN) Project CGL2009-12703-C03-03 and MICINN (BES-2010-030127) (R.R.-V.); Consejo Nacional de Ciencia y Tecnología (Mexico) (J.V.M.M.); Biotechnology and Biological Sciences Research Council BB/H005854/1 (V.W., F.B., and A.M.); European Research Council and Marie Curie Actions Grant 300554 (M.E.A.); Wenner-Gren Foundations and the Australian Research Council Future Fellowship FT0992258 (C.I.S.); European Research Council ERC-2011-AdG 295733 grant (Langelin) (D.P. and D.L.); Bernice Peltier Huber Charitable Trust (C.H. and L.G.D.); Russian Foundation for Basic Research grant 13-06-00670 (E.B.); Russian Foundation for Basic Research grant 14-0400725 (E.K.); European Union European Regional Development Fund through the Centre of Excellence in Genomics to Estonian Biocentre and Estonian Institutional Research grant IUT24-1 (E.M., K.T., M.M., M.K., and R.V.); Estonian Science Foundation grant 8973 (M.M.); Stanford Graduate Fellowship (J.R.H.); Washington State University (B.M.K.); French National Research Agency grant ANR-14-CE31-0013-01 (F.-X.R.); European Research Council grant 261213 (T.K.); National Science Foundation BCS-1025139 (R.S.M.); Social Science Research Council of Canada (K.-A.P. and V.G.); National Institutes of Health grants R01-GM094402 (M.S. and Y.S.S.), R01-AI17892 (P.J.N. and P.P.), and 2R01HG003229-09 (R.N. and C.D.B.); Packard Fellowship for Science and Engineering (Y.S.S.); Russian Science Fund grant 14-04-00827 and Presidium of Russian Academy of Sciences Molecular and Cell Biology Programme (O.B.); and Russian Foundation for Basic Research grant 14-06-00384 (Y.B.). Informed consent was obtained for the sequencing of the modern individuals, with ethical approval from the National Committee on Health Research Ethics, Denmark (H-3-2012-FSP21). SNP chip genotype data and whole-genome data for select present-day individuals are available only for demographic research under data access agreement with E.W. (a list of these samples is provided in tables S1 and S4). Raw reads from the ancient and the remainder of the present-day genomes are available for download through European Nucleotide Archive (ENA) accession no. PRJEB9733, and the corresponding alignment files are available at www.cbs.dtu.dk/suppl/NativeAmerican. The remainder of the SNP chip genotype data can be accessed through Gene Expression Omnibus (GEO) series accession no. GSE70987 and at www.ebc.ee/free_data. The authors declare no competing financial interests. C.D.B. is on the advisory board of Personalis, Identify Genomics, Etalon DX, and Ancestry.com.
- SUPPLEMENTARY MATERIALS**
- www.sciencemag.org/content/349/6250/aab3884/suppl/DC1
Materials and Methods
Supplementary Text
Figs. S1 to S41
Tables S1 to S15
References (65–205)
- 4 May 2015; accepted 15 July 2015
Published online 23 July 2015
[10.1126/science.aab3884](https://doi.org/10.1126/science.aab3884)

Genomic evidence for the Pleistocene and recent population history of Native Americans

Maanasa Raghavan, Matthias Steinrücken, Kelley Harris, Stephan Schiffels, Simon Rasmussen, Michael DeGiorgio, Anders Albrechtsen, Cristina Valdiosera, María C. Ávila-Arcos, Anna-Sapfo Malaspinas, Anders Eriksson, Ida Moltke, Mait Metspalu, Julian R. Homburger, Jeff Wall, Omar E. Cornejo, J. Víctor Moreno-Mayar, Thorfinn S. Korneliussen, Tracey Pierre, Morten Rasmussen, Paula F. Campos, Peter de Barros Damgaard, Morten E. Allentoft, John Lindo, Ene Metspalu, Ricardo Rodríguez-Varela, Josefina Mansilla, Celeste Henrickson, Andaine Seguin-Orlando, Helena Malmström, Thomas Stafford Jr., Suyash S. Shringarpure, Andrés Moreno-Estrada, Monika Karmin, Kristiina Tambets, Anders Bergström, Yali Xue, Vera Warmuth, Andrew D. Friend, Joy Singarayer, Paul Valdes, Francois Balloux, Ilán Lebreiro, Jose Luis Vera, Hector Rangel-Villalobos, Davide Pettener, Donata Luiselli, Loren G. Davis, Evelyne Heyer, Christoph P. E. Zollikofer, Marcia S. Ponce de León, Colin I. Smith, Vaughan Grimes, Kelly-Anne Pike, Michael Deal, Benjamin T. Fuller, Bernardo Arriaza, Vivien Standen, Maria F. Luz, Francois Ricaut, Niede Guidon, Ludmila Osipova, Mikhail I. Voevoda, Olga L. Posukh, Oleg Balanovsky, Maria Lavryashina, Yuri Bogunov, Elza Khusnutdinova, Marina Gubina, Elena Balanovska, Sardana Fedorova, Sergey Litvinov, Boris Malyarchuk, Miroslava Derenko, M. J. Mosher, David Archer, Jerome Cybulski, Barbara Petzelt, Joycelynn Mitchell, Rosita Worl, Paul J. Norman, Peter Parham, Brian M. Kemp, Toomas Kivisild, Chris Tyler-Smith, Manjinder S. Sandhu, Michael Crawford, Richard Villems, David Glenn Smith, Michael R. Waters, Ted Goebel, John R. Johnson, Ripan S. Malhi, Mattias Jakobsson, David J. Meltzer, Andrea Manica, Richard Durbin, Carlos D. Bustamante, Yun S. Song, Rasmus Nielsen and Eske Willerslev

Science **349** (6250), aab3884.

DOI: 10.1126/science.aab3884 originally published online July 21, 2015

Genetic history of Native Americans

Several theories have been put forth as to the origin and timing of when Native American ancestors entered the Americas. To clarify this controversy, Raghavan *et al.* examined the genomic variation among ancient and modern individuals from Asia and the Americas. There is no evidence for multiple waves of entry or recurrent gene flow with Asians in northern populations. The earliest migrations occurred no earlier than 23,000 years ago from Siberian ancestors. Amerindians and Athabascans originated from a single population, splitting approximately 13,000 years ago.

Science, this issue 10.1126/science.aab3884

ARTICLE TOOLS

<http://science.sciencemag.org/content/349/6250/aab3884>

SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2015/07/20/science.aab3884.DC1>

RELATED CONTENT

<http://science.sciencemag.org/content/sci/349/6246/358.full>
<http://science.sciencemag.org/content/sci/349/6246/354.full>
<http://science.sciencemag.org/content/sci/349/6246/372.full>
<http://science.sciencemag.org/content/sci/349/6246/370.full>
<http://science.sciencemag.org/content/sci/349/6246/367.full>
<http://science.sciencemag.org/content/sci/349/6246/441.2.full>
<http://science.sciencemag.org/content/sci/349/6246/362.full>
<http://science.sciencemag.org/content/sci/349/6246/369.full>
<file:/content/pending:yes>

Use of this article is subject to the [Terms of Service](#)

REFERENCES

This article cites 174 articles, 36 of which you can access for free
<http://science.sciencemag.org/content/349/6250/aab3884#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science* is a registered trademark of AAAS.